

Acanthamoeba Keratitis: Shriram H Bairagi*

Diagnosis and Treatment

Department of Pharmaceutical Chemistry,
Ideal College of Pharmacy and Research,
Mumbai, India

Received: July 13, 2017; Accepted: July 14, 2017; Published: July 19, 2017

*Corresponding author: Bairagi SH

✉ shrirambairagi@gmail.com

Assistant Professor, Department of
Pharmaceutical Chemistry, Ideal College of
Pharmacy and Research, Mumbai, 421306,
India.

Tel: +917715031328

Citation: Bairagi SH (2017) *Acanthamoeba*
Keratitis: Diagnosis and Treatment. J Rare
Disord Diagn Ther. Vol. 3 No. 2:8

Introduction

Acanthamoeba species are the causative agent of a sight threatening infection of the cornea known as *Acanthamoeba* keratitis, caused by the free-living amoeba *Acanthamoeba* spp., and is a sight-threatening corneal infection that presents with corneal ulcerations. It was first reported in 1973 [1]. *Acanthamoeba* are ubiquitous protozoa that exist in 2 forms: Trophozoites (the active form) and cysts (the inactive form). Cysts are notoriously difficult to kill and this is one reason why this infection is so difficult to eradicate. Only one class of medications is known to have cystocidal activity, the biguanides. The incidence of *Acanthamoeba* keratitis in this study was found to be 1.04% and the incidence in other reported Indian series ranges from 0.34 to 1.4% [2-5].

The incidence of *Acanthamoeba* keratitis appears to be increasing because of its frequent association with contact lens wear, which represents the cause of >85% of *Acanthamoeba* keratitis cases, especially in developed countries [6-8]. Members of the genus *Acanthamoeba* are ubiquitous and can be isolated from well, tap, bottled and swimming pool water, as well as sand, dust, human nasal and throat secretions, and animal stools. The life cycle of *Acanthamoeba* consists of a trophozoite and a cyst stage [9].

Diagnosis and Detection

The detection can be done in laboratory using non nutrient agar saline plate seeded with gram negative bacteria such as *Escherichia coli*. The PCR is used to confirm the diagnosis especially when the contact lenses are involved. Molecular methods also available for detection and identification of *Acanthamoeba*, these methods are also suitable for both clinical and epidemiological purposes [10,11]. The fluorescent *in situ* hybridization technique has also been successfully employed for the purpose detection [12]. **Figure 1** shows the infected eye.

Treatment and Case Studies

Acanthamoeba is difficult to treat, but effective management can save ones eye. Topical anti-infective agents used in early-stage, surgical intervention is necessary in later stages. Propamidine 0.1% and neomycin 1% shows good activity against AK Since then, two or more topical anti-amoebic agents are used throughout the day. Polyhexamethylene Biguanide (PHMB, 0.02%) and

chlorhexidine (0.02%), are effective in both monotherapy and combination therapy with a diamidine [13].

Case studies show that the *Acanthamoeba* keratitis is successfully treated in the person having the contact lens using six month




Figure 1 Infected eye.

therapy with topical Miconazole, Metronidazole, Prednisolone and neomycin as well as oral ketokonazole. In another case of person without contact lens having burning sensation the additional use of Propamidine in the above therapy shows the improvement in the corneal ulcer. In another case where topical Tobramycin and Cefazolin added in the therapy [14]. The combination of Dibromopropamidine and Propamidine isethionate ointment and drops and neomycin drops are used successfully in the 44 years old patient having corneal infection from *Acanthamoeba* species [15].

Bilateral *Acanthamoeba* keratitis was successfully treated with Chlorhexidine, Polyhexamethylene biguanide, Propamidine isethionate

in combination with Atropine and Dibromopropamidine an improvement was noted at which point Atropine and Dibromopropamidine were stopped while Fluorometholone acetate, Acyclovir and Ciprofloxacin were prescribed and patient was discharged [16]. Few studies state that addition of the neomycin-polymyxin B also gives the good result [17].

Prevention

Mostly the Contact lens wearers need to take precaution while cleaning the lenses should never use the tap water and saline to clean. As the treatment is toxic and lengthy so they should visit if they find out some sign of inflammation. And those without lenses they should not wash their eyes with lake or sea water.

References

- 1 Illingworth CD, Cook SD (1998) *Acanthamoeba* keratitis. *Surv Ophthalmol* 42: 493-508.
- 2 Manikandan P, Bhaskar M, Revathy R, John RK, Narendran V, et al. (2004) *Acanthamoeba* keratitis: A six year epidemiological review from a tertiary care eye hospital in South India. *Indian J Med Microbiol* 22: 226-230.
- 3 Srinivasan M, Gonzales CA, George C, Cevallos V, Mascarenhas JM, et al. (1997) Epidemiology and etiological diagnosis of corneal ulceration in Madurai, South India. *Br J Ophthalmol* 81: 965-971.
- 4 Leck AK, Thomas PA, Hagan M, Kalimurthy J, Ackuaku E, et al. Aetiology of suppurative corneal ulcers in Ghana and South India and epidemiology of fungal keratitis. *Br J Ophthalmol* 86: 1211-1215.
- 5 Basak SK, Basak S, Mohanta A, Bhowmick A (2005) Epidemiological and microbiological diagnosis of suppurative keratitis in Gangetic West Bengal, Eastern India. *Indian J Ophthalmol* 53: 17-22.
- 6 Patel A, Hammersmith K (2008) Contact lens-related microbial keratitis: recent outbreaks. *Curr Opin Ophthalmol* 19: 302-306.
- 7 Thebpatiphat N, Hammersmith KM, Rocha FN (2007) *Acanthamoeba* keratitis: a parasite on the rise. *Cornea* 26: 701-706.
- 8 Verani JR, Lorick SA, Yoder JS (2009) National outbreak of *Acanthamoeba* keratitis associated with use of a contact lens solution, United States. *Emerg Infect Dis* 15: 1236-1242.
- 9 Jonckheere DE (1991) JF: Ecology of *Acanthamoeba*. *Rev Infect Dis* 13: S385-S387.
- 10 Pasricha G, Sharma S, Garg P, Aggarwal R (2003) Use of 18S rRNA Gene-Based PCR Assay for Diagnosis of *Acanthamoeba* Keratitis in Non-Contact Lens Wearers in India *Journal of Clinical Microbiology* 41: 3206-3211.
- 11 Schroeder JM, Booton GC, Hay J, Niszl IA, Seal DV, et al. (2001) Use of subgenetic 18S ribosomal DNA PCR and sequencing for genus and genotype identification of *acanthamoebae* from humans with keratitis and from sewage sludge. *J Clin Microbiol* 39: 1903-1911.
- 12 Stothard DR, Hay J, Schroeder-Diedrich JM, Seal DV, Byers TJ (1999) Fluorescent oligonucleotide probes for clinical and environmental detection of *Acanthamoeba* and the T4 18S rRNA gene sequence type. *J Clin Microbiol* 37: 2687-2693.
- 13 Wright P, Warhurst D, Jones BR (1985) *Acanthamoeba* keratitis successfully treated medically. *Br J Ophthalmol* 69: 778-782.
- 14 Duff D, Horne MD, Mary EF (1994) *Acanthamoeba* keratitis: an emerging clinical problem. *Can Med Assoc J* 150: 1.
- 15 Wright P, Warhurst D, Barrie R (1985) *Acanthamoeba* keratitis successfully treated medically. *Br J Ophthalmol* 69: 778-782.
- 16 Hassanlou M, Bhargava A, Hodge WG (2006) *Acanthamoeba* keratitis and treatment strategy based on lesion depth. *Can J Ophthalmol* 41: 71-73.
- 17 Wang IJ, Hong JP, Hu FR (1997) Clinical features and outcome of *Acanthamoeba* keratitis. *J Formosan Med Assn* 96: 895-900.