

DOI: 10.21767/2380-7245.100036

Should All Abscesses be Drained? A Viral Infection of the Hand

Abstract

Orf is a zoonotic disease, which is caused by the parapox virus limiting itself and this disease can be transmitted by touching animals or infected flesh. This is usually seen on butchers and vets. The disease can be healed spontaneously and its treatment is symptomatic. The patients, going to hospitals complaining that there are abscesses on their hands, may encounter with wrong diagnosis and unnecessary surgical attempts. We are aimed at increasing the awareness of doctors to decide how to naje the treatment plans by taking the Orf on patients into consideration. On this study, we examined the Orf diagnosis, monitored on a 21-years old patient.

Keywords: Orf; Symptomatic treatment; Zoonoz; Parapox virus

Received: February 01, 2016; **Accepted:** February 09, 2016; **Published:** February 15, 2016

Introduction

Orf is a dermatologic disease known as contagious dermatitis or ecthyma. The disease is caused by the Orf virus and this virus is from the parapoxvirus family having double spiral [1]. It's also zoonoses that can be transmitted to human by touching animals like sheep and goat directly or infected flesh and contaminated waste. Characteristics Orf Virus Infections in Animals and Humans are shown in **Table 1**.

The risky group consists of farmers, butchers, vets and those distributing flesh. It has not been reported that it is transmitted from person to person [2]. The virus is seen as rare in the world and it is rarely reported to be seen as it limits itself [3]. The Orf, seen on human, is a solitary nodule which is 2-1 cm caliber. The lesion is generally seen on hands, wrists and forearms, but rarely on face. Clinically, in a week after the contact, there happens a red maculopapular lesion. This lesion is transformed into a nodule, on which there is a bulla. On this phase, lesion causes edema and it is so sensitive, bleeding very easily. In a week, there happens tiny pappilomes and these form thick crusts covering the surface. Then, it heals itself spontaneously in 6-8 weeks by giving out a superficial scar [1].

In recent years, there have been different alternative antiviral agents such as idoxuridine [3], cidofovir [4] and imiquimod [5] and shave excision cryotherapy [6] to heal the disease caused by Orf. We've found it suitable to present our phenomenon to examine Orf, which can be healed symptomatically without giving out any scars.

**Vahit Onur Gul¹,
Sebahattin Destek²,
Serhat Ozer³,
Serkan Ahioğlu¹ and
Yesim Erbil⁴**

- 1 Edremit Military Hospital General Surgery Department, 10300 Edremit/Balikesir, Turkey
- 2 Via Hospital General Surgery Department, 34947 Tuzla/Istanbul, Turkey
- 3 Konya Military Hospital General Surgery Department, 42040 Konya, Turkey
- 4 Istanbul University General Surgery Department, Istanbul, Turkey

Corresponding author: Gul VO

 vonurgul@hotmail.com

Edremit Military Hospital, General Surgery Department, Inonu St. 10300, Edremit Balikesir Turkey.

Tel: 905323669227

Citation: Gul VO, Destek S, Ozer S, et al. Should All Abscesses be Drained? A Viral Infection of the Hand. J Rare Dis Diagn Ther. 2016; 2:1.

Case

A 21 years-old man came to our clinic, complaining about lump and spasm on the first, 2nd, 4th fingers on his right hand and on surface of the same hand. He said he is a butcher and told us that he was in contact with zoic products. After the examination, we realized that there was hemorrhagic bulla which was erythematous and had edema on his right hand- first finger (0.5), right hand forth finger (0.5), and right hand third finger (0.5) and the nodules were so sensitive. On the right axilla, a lymph (1 cm) nod was detected.

On the right hand of the patient, there were widespread edema, hiperema and heat increase. On the liquid extracted from the lesion, it was realized that there were 2 neutrophil, on every tract. There was no reproduction culture. The diagnosis result was the Orf on the patient by taking clinic examination into consideration. In addition, the patient was given 1000 mg oral amoxicilin

clavulanate treatment, thinking there could be secondary infection. Having the patient used antibiotic for 10 days, it was detected the pain and edema were disappearing slowly. It was also monitored that the lesion disappeared spontaneously without any more intervention.

Discussion

The reason why the Orf disease is ignored much is because the disease limits itself [3]. For some time, it may lead to systemic fever and weakness. Especially, it is usual to see recurrence on risky people. On atopic dermatitis, transparent receivers and insufficient immune conditions, lesions may form exophytic masses, repeat themselves or become persistent [7]. The disease is generally seen in spring and summer months. Any sex separation hasn't been reported yet. All patients are males. Cause they are more in contact with animals [8]. The disease is generally seen on hands and faces, rarely on the other parts of the body [3]. The lesions emerge (**Figures 1 and 2**) as sole or a few on the hands which are not dominant [1].

But in our study, we realized that there was more than a single lesion on the dominant hand of the patient. In addition, Orf can be seen together with Stevens Johnson syndrome, generalized mucocutaneous eruption, edema on eyelid or big, stubborn and ongoing lesions on patients whose immune systems are too weak. On immunocompromised patients, there could be bigger

and fungal lesions [8]. On the other hand of our patient, we detected a cellulite with a widespread hyperemia and a sensitive lymphadenopathy (1 cm) on his armpit. After the oral amoksasilin clawlanate treatment, the lesions were healing spontaneously. It is easier to decide the diagnosis if we learn the patient is in contact with any animal. The lesions of our patient began with the contact with blood of a sheep in a week and the diagnosis was decided by taking his story and clinic studies into consideration. If someone wants to verify it, it can be done with the help of electron microscope, PCR or histopathological study. But histopathological study findings may change from patient to patient. These include pyknotic- epidemic cells, paleness on epidermis, eosinophilic inclusion, histiocytes, macrophages and lymphocyte infiltration [1]. In the distinctive findings of the disease, there may be anthrax, pyoderma, gangrenosum, herpetic felon, felon, milker's nodule malignant melanoma, staphylococcal dermatologic infections, atypical mycobacterial infections, tularemia and nokardiosis [9]. Specific antiviral treatment against the disease is not possible. On the same study, it was found that cidafovir, idoxuridine and imiquimod topical cream may give patients some benefits. In Orf treatment, cryotherapy, surgical excision and iterfenon were also used [3]. For the treatment of big lesions or on patients having weak immune systems, alfa interferon injection into lesions or cream with cidofovir can be used [10]. In addition, it slowly loses its population these days although 'shave' excision is a usual way to heal Orf disease and it is preferred by most doctors

Table 1 Characteristics of Orf virus infections in animals and humans.

	Animals	Humans
Incubation period	2–3 days	3–7 days
Mode of transmission	Direct contact	Direct contact
Clinical signs	Papules, pustules, vesicles	Papules that progress to pustules
Treatment	Supportive	Supportive
Course of disease	Self-limiting	Self-limiting
Complications	Anorexia, starvation, lameness, secondary bacterial infections	Secondary bacterial infections, bullous pemphigoid, erythema multiforme, toxic erythema
Vaccine	Yes	No
Important differential diagnoses	Foot and mouth disease	Cutaneous anthrax, tularemia, primary inoculation tuberculosis, eysipeloid, sporotrichosis



Figure 1 Lesions at first presentation.



Figure 2 Lesions after four weeks.

[6]. In Ireland, there is an attenuated vaccine and it is used for the Orf treatments [8]. It is also important to give importance to your hand hygiene and learn about the ways to protect yourself from this disease.

Conclusion

Today, there are not so many reports about this disease, but we may encounter with wrong treatment methods to heal the

disease. Mostly, surgical interventions rather than symptomatic are applied to heal patients complaining about scars and abscesses on their hands. Although this disease may disappear without applying to any treatment, there are lots of unnecessary surgical intervention and healing methods. With this study, we are aimed at increasing the awareness of doctors about arranging the true treatment plan by not ignoring patients' stories with animals.

References

- 1 Uzel M, Sasmaz S, Bakaris S, Cetinus E, Bilgic E, et al. (2005) A viral infection of the hand commonly seen after the feast of sacrifice: human orf (orf of the hand). *Epidemiol Infect* 133: 653-657.
- 2 Schmidt E, Weissbrich B, Brocker EB, Fleischer K, Goebeler M, et al. (2006) Orf followed by erythema multiforme. *J Eur Acad Dermatol Venereol* 20: 612-613.
- 3 Gurel MS, Ozardali I, Bitiren M, San I, Zeren H (2002) Giant orf on the nose. *Eur J Dermatol* 12: 183-185.
- 4 Damon IK (2010) Other poxviruses that infect humans: Parapoxviruses, Molluscum contagiosum, and Yatapoxviruses. In: *Principles and Practice of Infectious Diseases*. (7th edn) Philadelphia, Churchill Livingstone, Elsevier 1933-1936.
- 5 Erbagci Z, Erbagci I, Almila Tuncel A (2005) Rapid improvement of human orf (ecthyma contagiosum) with topical imiquimod cream: report of four complicated cases. *J Dermatolog Treat* 16: 353-356.
- 6 Depraeve C, De Coninck A, Senneseael J, Roseeuw D (1999) Recurrent contagious ecthyma (Orf) in an immunocompromised host successfully treated with cryotherapy. *Dermatology* 198: 162-63.
- 7 Ballanger F, Barbarot S, Mollat C, Bossard C, Cassagnau E, et al. (2006) Two giant orf lesions in a heart/lung transplant patient. *Eur J Dermatol* 16: 284-286.
- 8 Nadeem M, Curran P, Cooke R, Ryan CA, Connolly K (2010) Orf:contagious pustular dermatitis. *Ir Med J* 103: 152-153.
- 9 Frandsen J, Enslow M, Bowden AR (2011) Orf parapoxvirus infection from a cat scratch. *Dermatology Online Journal* 17: 9.
- 10 Geerinck K, Lukito G, Snoeck R, De Vos R, De Clercq E, et al. (2001) A case of human orf in a immunocompromised patient treated successfully with Cidofovir cream. *J Med Virol* 64: 543-549.