

Comparison of Surrogate Parameters between Cystic Fibrosis (CF) Patients in Frankfurt and Moscow during 1990-2015

Jean-Pascal Varescon^{1*},
Christina Smaczny¹,
Olaf Eickmeier^{1,3},
Gulja Babadjanova²,
Yulia Philippova²,
Stanislav Krasovski²,
Elena Amelina², and
Thomas Otto Friedrich
Wagner^{1,3}

Abstract

Background: Previous studies have demonstrated that CF (Cystic Fibrosis) prognosis is dependent of three major parameters: FEV1 (Forced Expiratory Pressure in one second), BMI (Body Mass Index) and need of intravenous antibiotic therapy. The CF centres of Frankfurt, Germany, and Moscow, Russia, care for cystic fibrosis patients. We decided to investigate and compare both centers from 1990 to 2015. No comparable study has been published so far.

Method: German patient data was collected from the national cystic fibrosis database "Muko.web". Missing values were extracted from the Hospital Information System. Russian patient data were taken directly from the medical records in Moscow. In a descriptive statistical analysis with Bias and R Studio the values were compared.

Result: A total of 428 patients from Moscow (217 male, 211 female; 348 (81,3%) were *P. aeruginosa* positive) and 159 patients from Frankfurt (92 male, 67 female; 137 (86,2%) with *P. aeruginosa* positive) were compared with regard to *P. aeruginosa* positivity, BMI, FEV1 and need of intravenous antibiotic therapy. CF patients in Moscow stratified by age groups had lower BMI than CF patients in Frankfurt (age 16-18: p=0,003; age 19-22: p=0,004; age 23-29: p<0,001; age 30-35: p<0,001; age 36-66: p=0,024). In a matching pairs analysis including 100 patients from Frankfurt and 100 patients from Moscow for the year 2015 FEV1 was significantly lower in Moscow patients (p<0,001).

Conclusion: BMI, FEV1 and need of intravenous therapy have significant impact on survival and on quality of life of CF patients. A lower BMI and a lower FEV1 result in a worse survival and determine the prognosis. This study showed a significant difference in prognostic parameters between Frankfurt and Moscow in the cross-sectional analysis for the year 2015. A further study should evaluate this difference to show whether this difference will be found over a longer period of time.

Keywords: Cystic fibrosis; BMI; FEV1; Intravenous Antibiotic Therapy; Lung Function; *P. Aeruginosa*; Surrogate Parameters

Received: January 13, 2021; **Accepted:** February 05, 2021; **Published:** February 12, 2021

Introduction

Cystic Fibrosis (CF) is a disease characterized by a loss of function of the Cystic Fibrosis Transmembrane Conductance Regulator (CFTR) in different organs [1]. Previous work has showed that prognosis in CF is related to Body-Mass-Index (BMI), Forced Expiratory Volume in 1 second (FEV1) and need of intravenous antibiotic therapy

[2], this is why these three parameters should be measured and monitored regularly. They have significant impact on survival and on quality of life of CF patients [3]. Disease progression in Cystic Fibrosis (CF) is marked by deterioration of a number of physiological systems [3] especially lung function is affected [2] progressively leading to pulmonary damage and in a final state to respiratory failure. The two centers of the Pulmonology Scientific

- 1 Christiane Herzog CF-Zentrum, Pneumologie – Klinikum der Goethe-Universität Frankfurt am Main, Germany
- 2 Pulmonology Scientific Research Institute – Lomonosov State University, Moscow
- 3 European Reference Network for Rare Diseases of the Respiratory System (ERN-LUNG); Frankfurt, Germany

*Corresponding author:

Jean-Pascal Varescon

✉ uni.j.varescon@laposte.net

Christiane Herzog CF-Zentrum, Pneumologie – Klinikum der Goethe-Universität Frankfurt am Main, Germany

Tel: +496963014232

Citation: Varescon JP, Smaczny C, Eickmeier O, Babadjanova G, Philippova Y, et al. (2021) Comparison of Surrogate Parameters between Cystic Fibrosis (CF) Patients in Frankfurt and Moscow during 1990-2015. J Rare Disord Diagn Ther Vol.7 No.1:2

Research Institute, Moscow and the University Hospital Frankfurt (Christiane Herzog CF-Zentrum) started a collaboration starting in July 2018. Both centers care for adult CF-patients, children as well as adults. A retrospective descriptive study was started to look for differences between patients treated in the Moscow CF center and the Frankfurt CF center from 1990 to 2015. If there was a significant and relevant difference this should be detected in a difference in the three surrogate markers of prognosis in CF: BMI [4], FEV1 [5] and the necessity of intravenous antibiotic therapy caused by exacerbations in CF [6]. BMI can be compared in stratified age classes. Higher BMI is related to better lung function test results (which improves consequently quality of life and survival) and in particular for underweight individuals a poorer prognosis has been reported [4]. FEV1 is the second parameter used to mark progression of CF lung disease progression and evaluate therapeutic efficacy [5]. Furthermore FEV1 is used as prognostic tool for mortality [7-9]. A third marker for the survival of CF patients is the need of intravenous antibiotic therapy as a result of a severe pulmonary exacerbation [6] or *P. aeruginosa* infection [10]. Exacerbations have a big consequence in terms of current morbidity as well as implications for long term morbidity and mortality [6-8]. *P. aeruginosa* presence is associated with faster rates of lung function decline in all age groups [10]. Need of intravenous antibiotic therapy consequently results in decreased survival. No comparable study has been published so far, i.e., there has been no published comparative descriptive study comparing CF patients in different settings in the past.

Materials and Methods

German patient data

German patient data were collected from the German national CF registry "muko.web" [11]. This registry was started 1995 under the name "Qualitätssicherung Mukoviszidose" has been renamed "muko.web". In the year 2015 ninety German CF centers took part in data gathering within Muko.web describing in much detail 5331 patients (median age 20; 56,5% adults; 51,8% men; 80 died in 2015; median dying age 32) [11]. Data collected from muko.web for the study were height, weight, BMI, FEV1, Forced Vital Capacity (FVC), year of birth, year of death, gender and *P. aeruginosa* presence. In addition to this date of diagnosis of *P. aeruginosa* infection of CF-patients in Frankfurt from 1990 to 2015 were retrieved. These data were anonymized and gathered into an Excel table. Missing values - in particular those describing the utilization of intravenous antibiotic therapy (not listed in muko.web) - were completed with data from the Hospital Medical record Information System (Orbis, Agfa) of the Frankfurt University Hospital. German patients were coded with the letter "f" and were associated to group 01. They received a three-digit numerical code "XXX".

Russian patient data

Russian patient data were collected directly from the medical files of the Pulmonology Scientific Research Institute, Moscow of the FMBA (Federal Medical-Biological Agency) of Russian Federation. Collected data were anonymized and regrouped in the same

standardized table as in Frankfurt. Russian patients were coded with the letter "m" and were associated to group 02. In the same way as in Frankfurt they received a three-digit numerical code "XXX".

Grouping of data

With this anonymized code data of both centers were aggregated in one data table. Gender information was coded with 01 for male patients and 02 for female patients. Body weight was expressed in kg (kilograms), body height in cm (centimeters), FEV1 in mL (milliliters), FVC in mL (milliliters). The presence of *P. aeruginosa* was coded with 01, the absence of *P. aeruginosa* with 02. Necessity of intravenous antibiotic treatment received the code 01 (02 coded not necessary intravenous antibiotic treatment).

Data analysis with Bias and RStudio

The complete data were biometrically analyzed with the program "Bias" [12]. After a descriptive approach of complete data, differences in BMI and FEV1 values were observed. Exemplarily BMI in the two centers was compared for 2015 after separating the sample in age classes [13,14]. FEV1 is an inconstant value, as it depends on age, height and sex category [15,16]. To compare FEV1 between both centers in 2015 a program was written in Rcode and executed with RStudio –a statistical programming tool, which can execute Rcode and analyze statistical data-. MatchIt [17] was used to create two new comparable samples. They were matched according to the parameters influencing FEV1 (height, age, sex category) [15,16]. The size of both samples was 100 patients and both samples were statistically not significant different (before matching $p < 0,001$, after matching $p = 0,484$). Afterward both new samples were compared for their FEV1 values in a new statistical biometrical analysis with "Bias" [12].

Comparison to normal population

At the end the descriptive data of the study were compared to normal population data in Germany [18] and the Russian Federation [19-21] taking in consideration epidemiological differences, which can influence CF-patients in both centers. Especially differences in BMI in both countries may have an influence on samples BMI.

Results

Description of the data range: Year of birth

The study totalized 428 (72,91%) patients from Moscow and 159 (27,09%) patients from Frankfurt, what conducted to a total of 587 analyzed patients. All of them were born in 1999 or earlier. The oldest patient of this study was born in 1949 (Figure 1). Average [22] year of birth was 1982 for Frankfurt, 1988 for Moscow and for the total cohort 1986. Standard deviation [22] was 11,10 for Frankfurt, 6,42 for Moscow and 8,45 for total cohort. Median [22] year of birth was 1983 for Frankfurt (1st quartile 1973, 3rd quartile 1992), 1989 for Moscow (1st quartile 1985, 3rd quartile 1993) and 1988 for total cohort (1st quartile 1983, 3rd quartile 1993). Minimum in Frankfurt was 1956. In Moscow it was 1949 (consequently 1949 for total cohort). In both centers maximum




Figure 1 Differences in year of birth in both CF-populations. (A) Year of birth distribution of CF-patients in both centers described in dot plots. (B) Box plot of patients year of birth representing median, first quartile and third quartile. (C) Box plot of patient's year of birth representing average and standard deviation.

was 1999 (so same for total cohort. Finally range was 43 for Frankfurt and 50 for Moscow and total cohort.

Description of the data range: Gender distribution, *P. aeruginosa* presence and more

In Moscow 217 (50,70%) male patients and 211 (49,30%) female patients were counted, while in Frankfurt 92 (57,86%) male patients and 67 (42,14%) female patients were registered. A performed Chi-square test with Yates's correction for continuity, confirms both samples were comparable ($p=0,147$) for sex category distribution. 348 (81,31%) Moscow patients were infected with *P. aeruginosa* and 80 (18,69%) were not infected with this bacterium. In Frankfurt the number of *P. aeruginosa* infected patients was 137 (86,16%) while 22 CF patients were not infected (13,84%). In the same way as for the sex category distribution a Chi-square test with Yates's correction for continuity [22] was performed ($p=0,209$). Subsequently both CF-patient populations were comparable for *P. aeruginosa* infections ($p=0,209$). 6 (3,77%) recorded patients died in Frankfurt (0 until 2015) and 114 (26,64%) recorded patients died in Moscow (68 (15,89%) until 2015).

Description of the data range: BMI, FEV1 and Necessity of intravenous antibiotic therapy evolution over time

Data were statistically analyzed and important values were calculated and entered into **Table 1**. Key values of BMI, FEV1

Table 1 (A) BMI biometrical descriptive statistic from 1990 to 2015 including number of patient data sets, average BMI, median BMI, SD (standard deviation) BMI, maximum BMI, minimum BMI, BMI range, 1st quartile BMI and 3rd quartile BMI. **(B)** FEV1 biometrical descriptive statistic from 1990 to 2015 including number of patient data, average FEV1, median FEV1, SD (standard deviation) FEV1, maximum FEV1, minimum FEV1, FEV1 range, 1st quartile FEV1 and 3rd quartile FEV1. **(C)** Biometrical descriptive statistical analysis of necessity of intravenous antibiotic therapy from 1990 to 2015 including number of patient data, number of necessity of intravenous antibiotic therapy and percentage of necessity of intravenous antibiotic therapy.

A)

Year	Number of Patient data		BMI average		BMI median	
	Frankfurt	Moscow	Frankfurt	Moscow	Frankfurt	Moscow
1990	2	0	21,52	-	21,52	-
1991	2	1	21,54	14,49	21,54	14,49
1992	1	2	21,50	16,27	21,50	16,27
1993	2	7	20,91	16,44	20,91	15,08
1994	1	11	20,02	17,34	20,02	17,16
1995	10	7	16,66	17,36	16,11	17,16
1996	30	10	18,84	16,91	18,52	17,39
1997	35	23	19,41	17,61	19,55	17,72
1998	46	38	19,89	16,34	19,66	16,45
1999	45	39	19,97	17,33	19,13	17,65
2000	30	45	20,29	16,98	19,09	16,53
2001	14	53	19,07	16,79	18,05	16,85
2002	16	64	18,78	17,41	18,02	17,54
2003	68	78	21,43	17,42	20,85	17,55
2004	75	103	21,48	17,49	20,76	17,57
2005	13	101	20,57	18,04	20,68	18,03
2006	13	124	20,95	18,02	21,27	17,96
2007	13	160	20,51	18,11	21,10	18,13
2008	91	179	21,69	18,38	21,01	18,55
2009	84	188	22,39	18,55	21,81	18,52
2010	132	192	21,20	18,76	20,70	18,69
2011	137	199	21,40	18,71	21,14	18,47
2012	131	250	21,80	18,79	21,62	18,51
2013	130	263	21,99	18,68	21,66	18,29
2014	133	278	22,12	18,78	21,73	18,52
2015	141	301	22,24	18,74	21,63	18,59
Year	BMI standard deviation (SD)		BMI maximum		BMI minimum	
	Frankfurt	Moscow	Frankfurt	Moscow	Frankfurt	Moscow
1990	0,64	-	21,98	-	21,07	-
1991	1,13	-	22,34	14,49	20,75	14,49
1992	-	1,75	21,50	17,51	21,50	15,03
1993	1,26	2,88	21,80	20,93	20,02	13,22
1994	-	3,82	20,02	25,00	20,02	13,34
1995	2,29	2,37	20,64	22,21	13,68	15,43

Year	Number of Patient data		BMI average		BMI median	
	Frankfurt	Moscow	Frankfurt	Moscow	Frankfurt	Moscow
1996	3,17	2,21	25,83	19,37	14,07	13,47
1997	3,18	2,49	28,22	22,77	13,71	13,34
1998	3,18	3,58	27,64	22,77	13,65	1,92
1999	3,52	2,51	31,11	22,94	14,88	12,63
2000	4,89	2,71	37,56	24,15	13,13	12,70
2001	4,02	2,69	27,76	22,76	14,60	12,40
2002	2,94	2,65	24,01	23,23	14,74	11,65
2003	3,76	2,70	33,30	23,61	13,98	11,65
2004	4,05	2,77	35,50	24,88	12,93	10,82
2005	3,22	2,76	26,35	25,86	13,73	12,02
2006	3,35	2,83	26,67	25,72	14,38	12,03
2007	3,58	2,73	24,97	25,62	13,89	12,73
2008	4,10	2,79	40,75	26,23	14,38	11,83
2009	4,41	2,78	44,29	26,03	15,34	12,80
2010	4,20	2,84	45,35	30,03	14,27	12,60
2011	4,18	2,90	45,52	31,99	14,35	12,47
2012	4,03	2,69	45,34	27,73	13,86	12,47
2013	4,04	2,80	44,47	27,73	13,86	10,85
2014	4,17	2,86	45,41	31,46	14,10	13,02
2015	4,13	2,78	46,60	31,46	14,17	11,33
Year	BMI range		BMI 1 st quartile		BMI 3 rd quartile	
	Frankfurt	Moscow	Frankfurt	Moscow	Frankfurt	Moscow
1990	0,91	-	-	-	-	-
1991	1,59	0,00	-	-	-	-
1992	0,00	2,48	-	-	-	-
1993	1,78	7,71	-	14,49	-	18,42
1994	0,00	11,66	-	14,22	-	19,30
1995	6,96	6,78	15,21	15,64	17,29	17,70
1996	11,76	5,90	16,51	15,35	20,88	18,85
1997	14,51	9,43	17,38	15,89	20,87	19,12
1998	13,99	20,85	17,93	14,22	21,60	18,46
1999	16,23	10,32	17,79	15,41	21,72	19,00
2000	24,43	11,45	17,59	14,81	21,15	19,23
2001	13,16	10,36	15,90	14,66	21,14	18,67
2002	9,27	11,58	16,76	15,23	21,38	18,93
2003	19,33	11,96	19,03	15,23	22,92	19,11
2004	22,57	14,06	19,23	15,21	23,00	19,47
2005	12,61	13,85	19,33	16,37	22,01	19,68
2006	12,29	13,69	19,76	15,66	22,60	19,82
2007	11,08	12,89	20,48	16,28	22,92	19,91
2008	26,37	14,40	19,58	16,47	22,80	20,09
2009	28,95	13,22	20,03	16,71	23,46	19,93
2010	31,08	17,43	18,81	16,97	22,95	20,20
2011	31,17	19,53	19,05	16,93	23,13	20,45
2012	31,48	15,27	19,34	16,86	23,29	20,45
2013	30,61	16,88	19,58	16,82	23,69	20,43
2014	31,32	18,44	19,31	16,86	23,81	20,50
2015	32,43	20,13	19,31	16,82	24,14	20,32

and necessity of intravenous antibiotic therapy were examined. Data were statistically analyzed and values were calculated and entered into **Table 1**. Key values of BMI, FEV1 and necessity of intravenous antibiotic therapy were examined. In summary the

parameters in **Table 1** are mostly better for Frankfurt patients than for Moscow patients. From 1990 to 1995 there were not sufficient data and consequently values and results cannot be considered to be representative. FEV1 values have to corrected by height, age and sex category [15,16], therefore, they are not directly comparable.

Statistical BMI comparison of both CF populations in 2015

To evaluate if there was a significant statistical BMI difference between patients in Frankfurt and Moscow year 2015 was analyzed exemplarily. Patients were categorized in age groups (**Figures 2 and 3**). In 2015 Moscow CF patients stratified by age groups had statistically significant lower BMI than Frankfurt CF patients in all age groups (age 16-18: p=0,003; age 19-22: p=0,004; age 23-29: p<0,001; age 30-35: p<0,001; age 36-66: p=0,024) [22-25].




Figure 2 Box plots representing BMI comparison of patients for 2015 with median, first quartile and third quartile. **(A)** Patients aged 16 to 18 years (Average BMI: Frankfurt (n=12): 19, 95; Moscow (n=25): 17, 90). **(B)** Patients aged 19 to 22 years (Average BMI: Frankfurt (n=19): 20, 87; Moscow (n=76): 18, 75). **(C)** Patients aged 23 to 29 years (Average BMI: Frankfurt (n=25): 22, 59; Moscow (n=139): 18, 66).




Figure 3 Box plots representing BMI comparison of patients for 2015 with median, first quartile and third quartile. **(A)** Patients aged 30 to 35 years (Average BMI: Frankfurt (n=29): 22, 27; Moscow (n=44): 18, 19). **(B)** Patients older than 35 years (Average BMI: Frankfurt (n=56): 23, 03; Moscow (n=17): 19, 93).

Statistical FEV1 comparison of matched samples in 2015

To compare FEV1 in both centers a program run with RStudio [17] allowed isolation of two matched samples by height (before matching $p=0,028$, after matching $p=0,876$), age (before matching $p<0,001$, after matching $p=0,484$) and sex category (before matching $p=0,088$, after matching $p=0,258$) for 2015. Both included 100 patients (first sample with 100 Frankfurt patients and second sample with 100 Moscow patients) and were comparable after matching. Statistical analysis showed FEV1 was significantly lower for Moscow CF-patients ($p<0,001$) than for Frankfurt CF-patients in 2015 (Table 2 and Figure 4).

Discussion

Data described both CF-populations in Frankfurt and Moscow. At first glance values of BMI, FEV1 and the necessity of intravenous antibiotic therapy were better in Frankfurt than in Moscow. An evaluation of both CF-populations for 2015 revealed BMI was significantly higher in Frankfurt, than in Moscow. A high BMI is a positive predictor for a better outcome [26,27] and decreased mortality [26]. Epidemiological analysis of normal German [18] and Russian [19-21] population didn't explain this severe gap (referred to 3.6.). In the same way both FEV1 populations of 2015 obtained with the R-program to get comparable samples showed Frankfurt CF-patients have a better FEV1 than Moscow CF-patients. A better FEV1 is associated with a better outcome [26] and a lower mortality. Subsequently these data indicate

Frankfurt patients should have a better outcome than Moscow patients. An effort to increase BMI and FEV1 will improve the CF prognosis in Moscow.

Comparison with BMI and FEV1 of normal population

At first we had to analyze epidemiological available data of normal population to see if both are reasonably comparable. Latest data from the German federal office of statistics [28] shows a mean BMI of 26.0 for German population in 2017. Russian data [29,30] are not equally detailed and latest data was published in 2014. Mean BMI in the Russian population was 26.5. In the same year mean BMI was 26.3 in Germany, this might mean that the Russian population has a higher mean BMI than the German population, however both populations can be considered comparable. Consequently a possible gap in BMI in both CF-populations (referred to 3.4.) cannot be explained by epidemiological data of the normal population. A comparison of FEV1% between Germany and the Russian Federation [31] shows a difference for patients categorized in age groups. Average and mean values seem higher in Germany for children and for adults. According to ECFSPR annual report of 2017 [31] the FEV1% of Germany and the Russian Federation are different. German data seem to resemble the pooled data very closely, while the Russian data seem to be lower than pooled data and German data. This is the reason why we expected differences between both centers we wanted to analyze.

Limitations of the study


The data quality of our study should be discussed. First of all, it should be mentioned that data were not available from every patient every year. This is why the significance of the data should be nuanced. As an example, in 2015 for the entire cohort, only 486 out of 587 entries (82.79%) were found for the BMI, only 440 out of 587 (74.96%) entries were found for FEV1 and only 492 out of 587 (83.82%) entries were found for intravenous antibiotic therapy. This shows that a significant amount of data is missing and that the quality of the data is negatively affected. Moreover, the data was collected on one hand by doctors and clinic employees, which makes human bias in the data collection possible. On the other hand, this clinically collected data is entered manually into the computer system, which makes further errors possible and can explain missing data. Deviations due to anomalies were also found in the patient's follow-up data. These have also affected the quality of the data and thus reduced the representativeness of the data.

Possible explanation for the observed differences

In our study we could observe the Russian cohort is significantly younger than the German cohort. Average age was 33.57 for Frankfurt, 25.59 for Moscow and for the total cohort 28.14. According to ECFSPR in 2017 mean average age was 22.4 [31] years in Germany and 12.4 [31] years in Russian Federation what confirms our results. In Moscow 217 (50.70%) male patients and 211 (49.30%) female patients were counted, while in Frankfurt

Table 2 Biometrical statistic analysis of FEV1 in 2015 for both samples (n=100 CF-patients in Frankfurt and n=100 CF-patients in Moscow). Average, median, SD, maximum, minimum, range, first quartile and third quartile are higher in Frankfurt than in Moscow.

FEV1	Average	Median	SD	Maximum	Minimum	Range	1 st quartile	3 rd quartile
Frankfurt	2497,90	2420,00	1143,19	5410,00	750,00	4660,00	1537,50	3325,00
Moscow	1908,70	1615,00	1044,80	5220,00	520,00	4700,00	1222,50	2487,50



92 (57.86%) male patients and 67 (42.14%) female patients were registered. According to ECFSPR in 2017 in Germany around 52% were male patients and 48% were female. In Russian Federation the percentage was similar with about 51% male patients and 49% female patients [31]. These results were comparable with our study for Moscow. In Frankfurt the relative amount of male patients was higher than the German average. The observed gender gap could have influenced our results.

According to the number of death patient totalized in our study, there were less deaths in Frankfurt than in Moscow. This can be partially explained by new therapies [32], a better organization [33] and a medicine that becomes more and more detailed and precise due to the economic possibilities and the research. This means that German patients are in a transition phase, where life expectancy increases. Patients in Russia yet are not in this phase. This may be linked to a possible delay in the use of more modern equipment and therapies, as well as probably lower or unevenly distributed financial means. Moreover the economic structures are different between both countries as well as the regional structures of Frankfurt and Moscow. Russia evaluated recently with the independence from the Soviet Union in 1991 and inherited an extensive centralized system. In 1993 a mandatory health insurance (MHI) was introduced to open up an earmarked stream of funding for health care, but faced

lots of fiscal constraints [34]. In Germany, the health system is build up in a different way. The state is organized federally and multiple adapted health care centers were created. This was also reflected in CF management. Since 1995, the German Cystic Fibrosis Quality Assessment project has collected demographic data and outcome parameters, what aims to develop tools for quality management and improve health care [35]. This could also partly explain our results, but has to be confirmed in further studies, where economical, management and organization can be monitored.

Important and new aspect of our study

Our study is the first one comparing CF patients between both centers of Frankfurt and Moscow. It has confirmed expected differences between surrogate parameters of prognosis (BMI, FEV1 and need of intravenous therapy) in CF patients of both centers. A higher BMI and FEV1 in one CF centre (Frankfurt) are positive prognostic parameters for survival compared to the values in Moscow. An effort to increase BMI and FEV1 in Moscow will certainly improve lifetime prognosis. This opens new ways to research possible causes of BMI gaps and FEV1 gaps and to close them. Our data indicates that in daily CF therapy routine BMI should be checked more often. Moreover higher BMI values should be targeted in CF patients. In a same way FEV1 should be monitored even more regularly and control intervals should be reduced. Our results permit to evaluate differences in therapy schemes and the use of various CF medications, in particular CFTR modulator therapies in further studies. Moreover our results show that the necessity of intravenous antibiotic treatment has also to be reexamined. The observed better results in Frankfurt for intravenous therapies in CF patients have to be proved statistically. The relation between the necessity of intravenous antibiotic therapy and a worse outcome for CF-patients has to be discussed. A study published in 2015 questioned the link between both and put other antibiotic treatments (oral therapy or inhaled therapy) on the same acting level [36]. Our work will also revolutionize research, as the focus of comparative and new founding studies should concentrate on the BMI, the FEV1 and the need of intravenous therapies. Furthermore regional development aspects and organizational differences have to be included and considered more often. The health-policy could also navigate on these findings to elaborate new health plans and new goal achievements based in particular on a higher BMI, a higher FEV1 and an evaluation of antibiotic use depending on the region and the available resources. Furthermore, we showed the reasons of these gaps in surrogate markers for CF prognosis have to be investigated. One possible cause could be a possible difference in delta F508 mutation or other CF-specific gene mutation distribution. Moreover epidemiologic reasons should be regarded in a larger scale and also compared and evaluated

in another study. The socioeconomic differences between both countries should also be taken in consideration. Our study indicates in particular that different types of drugs, modes of application, frequency of application, treatment regimens and the availability of medication could play a role in CF prognosis.

Conclusion

We have identified that Frankfurt CF patients values for surrogate parameters of CF outcome were better than those in Moscow patients in a short time. Further studies should verify this difference on a longer lapse of time including larger data spectrum. First, this will allow to establish a hypothesis explaining this difference. Secondly, this could help to refine therapeutic approaches and to definite new recommendations.

Acknowledgement

This research was supported by the Christiane Herzog CF-Zentrum, Pneumologie – Klinikum der Goethe-Universität Frankfurt am Main – and the Pulmonology Scientific Research Institute – Lomonosov State University Moscow –. We are thankful to our colleagues who provided expertise that greatly assisted the research. We are also grateful to Dr. Ümniye Balaban for assistance with a statistical help, and Prof. Dr. med. Gernot Rohde who moderated this paper and in that line improved the manuscript significantly.

Statement of Ethics

There are no legal or ethical concerns. The ethics committee issued a positive assessment. See document “Ethical approval Varescon Study”. Study number 183/17.

Conflict of Interest

The authors have no conflicts of interest to declare.

Funding Sources

None.

Author Contributions

Jean-Pascal Varescon: Conceptualization, data curation, methodology, software, formal analysis, investigation, resources, writing, review – original draft, visualization, project administration. Approval for submission, agreement to be

accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

T.O.F. Wagner: Supervision, conceptualization, writing – review & editing, project administration, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Gulja Babadjanova: Resources acquisition, Conceptualization, Methodology, review, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Christina Smaczny: Methodology and analysis, Resources acquisition, Conceptualization, review, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Olaf Eickmeier: Resources, review, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Yulia Philippova: Data curation, Resources, review, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Stanislav Krasovskiy: Resources, Data curation, review, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Elena Amelina: Resources, review, final approval for submission, agreement to be accountable for all aspects of the work is ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated or resolved.

Prof. Dr. med. Gernot Rohde: Substantial contributions to the conception of the work.

Dr. Ümniye Balaban: Statistical analysis help.

References

- 1 Meng X, Clews J, Ciuta AD, Martin ER, Ford RC, et al. (2019) CFTR structure, stability, function and regulation. *Biol Chem* 400: 1359-1370.
- 2 Kerem E, Conway S, Elborn S, Heijerman H (2005) Standards of care for patients with cystic fibrosis: a European consensus. *J Cyst Fibros* 4: 7-26.
- 3 Gee L, Abbott J, Conway SP, Etherington C, Webb AK, et al. (2003) Quality of life in cystic fibrosis: the impact of gender, general health perceptions and disease severity. *J Cyst Fibros* 2: 206-213.
- 4 Stephenson AL, Mannik LA, Walsh S, Brotherwood M, Robert R, et al. (2013) Longitudinal trends in nutritional status and the relation between lung function and BMI in cystic fibrosis: a population-based cohort study. *Am J Clin Nutr* 97: 872-877.
- 5 Szczesniak R, Heltshe SL, Stanojevic S, Mayer-Hamblett N (2017) Use of FEV1 in cystic fibrosis epidemiologic studies and clinical trials: A statistical perspective for the clinical researcher. *J Cyst Fibros* 16: 318-326.
- 6 Bhatt JM (2013) Treatment of pulmonary exacerbations in cystic fibrosis. *Eur Respir Rev* 22: 205-216.

- 7 Hayllar KM, Williams SG, Wise AE, Pouria S, Lombard M, et al. (1997) A prognostic model for the prediction of survival in cystic fibrosis. *Thorax* 52: 313-317.
- 8 Liou TG, Adler FR, Fitzsimmons SC, Cahill BC, Hibbs JR, et al. (2001) Predictive 5-year survivorship model of cystic fibrosis. *Am J Epidemiol* 153: 345-352.
- 9 Kerem E, Reisman J, Corey M, Canny GJ, Levison H, et al. (1992) Prediction of mortality in patients with cystic fibrosis. *N Engl J Med* 326: 1187-1191.
- 10 Keating C, Poor AD, Liu X, Chiuzan C, Backenroth D, et al. (2017) Reduced survival in adult cystic fibrosis despite attenuated lung function decline. *J Cyst Fibros* 16: 78-84.
- 11 Nährlich L, Burkhardt M, Wiese B (2015) German CF-Registry Annual Report 2015.
- 12 Ackermann H (2006) BiAS: Biometrische Analyse von Stichproben. 8th ed. Frankfurt am M.
- 13 Yi S-W, Ohrr H, Shin S-A, Yi J-J (2015) Sex-age-specific association of body mass index with all-cause mortality among 12.8 million Korean adults: A prospective cohort study. *Int J Epidemiol* 44: 1696-1705.
- 14 Hayes A, Gearon E, Backholer K, Bauman A, Peeters A (2016) Age-specific changes in BMI and BMI distribution among Australian adults using cross-sectional surveys from 1980 to 2008. *Int J Obes* 39: 1209-1216.
- 15 Lungenfunktionstest - Normwerte | Leichter-atmen.de. <https://www.leichter-atmen.de/lungenfunktionstest-werte>
- 16 Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, et al. (2012) Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *Eur Respir J* 40: 1324-1343.
- 17 Ho DE, Imai K, King G, Stuart EA (2011) MatchIt nonparametric preprocessing for parametric causal inference. *J Stat Soft* 42.
- 18 Statistisches Bundesamt der Bundesrepublik Deutschland. Körpermaße nach Altersgruppen und Geschlecht. <https://www.destatis.de/DE/Themen/Gesellschaft-Umwelt/Gesundheit/Gesundheitszustand-Relevantes-Verhalten/Tabellen/liste-koerpermasse.html;jsessionid=C7F5C7ACF13C6978E29DDF5BECA2E9C1.internet732>.
- 19 Shalnova S, Vilkov V, Balanova Y, Muromtseva G, Kapustina A, et al. (2018) P4451 Comparison of the body mass index in the populations of the Russian Federation and the United States of America during thirty years period. *Eur Heart J* 39.
- 20 World Health Organization (2013) Regional Office for Europe. Russian Federation - WHO Country Profile.
- 21 Rtveladze K, Marsh T, Webber L, Kilpi F, Goryakin Y, et al. (2012) Obesity trends in Russia. The impact on health and healthcare costs. *Health* 04: 1471-1484.
- 22 Sachs L (2004) *Angewandte Statistik*. Springer: Berlin Heidelberg.
- 23 Toutenburg H (1975) Hollander, M., D. A. Wolfe: *Nonparametric statistical methods*. John Wiley & Sons, New York-Sydney-Tokyo-Mexico City 1973. 503 S., \$9.50. *Biom J* 17: 526.
- 24 Cohen J (2013) *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hoboken: Taylor and Francis.
- 25 Zimmermann H (1985) Exact Calculation of Permutational Distributions for Two Independent Samples. *Biom J* 27: 431-434.
- 26 Corey M, McLaughlin FJ, Williams M, Levison H (1988) A comparison of survival, growth, and pulmonary function in patients with cystic fibrosis in Boston and Toronto. *J Clin Epidemiol* 41: 583-591.
- 27 Sharma R, Florea VG, Bolger AP, Doehner W, Florea ND, et al. (2001) Wasting as an independent predictor of mortality in patients with cystic fibrosis. *Thorax* 56: 746-750.
- 28 Statistisches Bundesamt der Bundesrepublik Deutschland. Körpermaße nach Altersgruppen und Geschlecht. <https://www.destatis.de/DE/Themen/Gesellschaft-Umwelt/Gesundheit/Gesundheitszustand-Relevantes-Verhalten/Tabellen/liste-koerpermasse.html;jsessionid=C7F5C7ACF13C6978E29DDF5BECA2E9C1.internet732>.
- 29 World Health Organization (2019) Regional Office for Europe. Russian Federation - WHO Country Profile.
- 30 Rtveladze K, Marsh T, Webber L, et al. Obesity trends in Russia. The impact on health and healthcare costs. *Health*. 2012;04(12):1471-1484.
- 31 European Cystic Fibrosis Society (2017) ECFS Patient Registry - Annual Data Report 2017.
- 32 Lopes-Pacheco M (2019) CFTR Modulators: The Changing Face of Cystic Fibrosis in the Era of Precision Medicine. *Front Pharmacol* 10: 1662.
- 33 Smaczny C, Eickmeier O, Wagner TOF (2013) Transition in der Pneumologie. *Pneumologie*. 10: 13-19.
- 34 Popovich L, Potapchik E, Shishkin S, Richardson E, Vacroux A, et al. (2011) Russian Federation. Health system review. *Health Syst Transit* 13: 1-190.
- 35 Stern M, Wiedemann B, Wenzlaff P (2006) From registry to quality management: the German Cystic Fibrosis Quality Assessment project 1995 2006. *Eur Respir J*. 2008;31: 29-35.
- 36 Hurley MN, Prayle AP, Flume P (2015) Intravenous antibiotics for pulmonary exacerbations in people with cystic fibrosis. *Cochrane Database Syst Rev* 2015: CD009730.